

Exact Result for the Nonlocal Conserved Kardar-Parisi-Zhang Equation

Eytan Katzav*

*School of Physics and Astronomy, Raymond and Beverly Sackler Faculty of Exact Sciences,
Tel Aviv University, Tel Aviv 69978, Israel*

Abstract

I analyze the Nonlocal Conserved Kardar-Parisi-Zhang (NCKPZ) equation with spatially correlated noise. This equation is also known as the Nonlocal Molecular Beam Epitaxy (NMBE) equation and was originally suggested to study the effect of the long-range nature of interactions coupled with spatially correlated noise on the dynamics of a volume conserving surface. I find an exact result for a subfamily of NCKPZ models in one dimension, and thus establish an exact result for MBE processes for the first time. Then, to complete the picture, I construct a Self-Consistent Expansion (SCE) and get results that are consistent with the exact result in one dimension. I conclude by discussing previous results obtained for NCKPZ using dynamic renormalization group approach, and find that this approach does not yield the exact result I obtain. This discussion shows the advantage of the using SCE to deal with non-linear stochastic equations.

*Electronic address: eytak@post.tau.ac.il

The field of disorderly surface growth has received much attention during the last two decades. A multitude of different phenomena such as fluid flow in porous media, propagation of flame fronts, flux lines in superconductors not to mention deposition processes, bacterial growth and "DNA walk" [1] are all said to be related to the famous Kardar-Parisi-Zhang (KPZ) equation [2].

In spite of the great success of KPZ to describe many growth models and phenomena, there has been a growing pool of data that is not well described by KPZ, that called for further research. One of the first classes that belongs to this non KPZ behavior is the well known conserved KPZ equation class (sometimes also called the Molecular-Beam-Epitaxy (MBE) class) [3]-[9]. However, the modified behavior introduced by the MBE equation is not at all sufficient to account for all the rich non KPZ experimental data in the field.

A different line of research suggested that the basic growth equation should not be changed. Instead, the white noise that appears in the original KPZ equation should be either temporally or spatially correlated [10]-[12]. This approach was indeed quite successful, but it still failed to give a good account for all the measured scaling exponents.

Recently, Mukherji and Bhattacharjee [13] proposed a phenomenological equation that takes into account the presence of long-range interactions. This equation, known as the Nonlocal KPZ (NKPZ) equation, has a nonlinear term that couples the gradients of the growing surface at two different points. This equation was studied using the Dynamic Renormalization Group (DRG) method [13] and so was its generalization that includes spatially correlated noise [14]. It turned out that a very rich family of phases was found.

Just like in the local case, a nonlocal conserved KPZ (NCKPZ) equation was needed in order to account for the dynamics of a volume conserving surface that experiences long-range interactions. This equation was proposed by Jung *et al.* [15, 16]. The continuum equation they suggest for the height of the surface $h(\vec{r}, t)$ at a point $\vec{r}$ and time t measured relative to its spatial average is

$$\begin{aligned} \frac{\partial h(\vec{r}, t)}{\partial t} = & -K\nabla^4 h(\vec{r}, t) - \frac{1}{2}\nabla^2 \int d^d r' g(\vec{r}') \\ & \times \nabla h(\vec{r} + \vec{r}', t) \cdot \nabla h(\vec{r} - \vec{r}', t) + \eta(\vec{r}, t), \end{aligned} \quad (1)$$

where K is a constant, the kernel $g(\vec{r})$ represents the long-range interactions with a short-range part $\lambda_0 \delta^d(\vec{r})$ and a long-range part $\sim \lambda_\rho r^{\rho-d}$, where d is the substrate dimension and ρ is an exponent characterizing the decay of the long range interaction. More precisely in

Fourier space $\hat{g}(q) = \lambda_0 + \lambda_\rho q^{-\rho}$. Finally, $\eta(\vec{r}, t)$ is a conservative spatially correlated noise term that satisfies

$$\begin{aligned}\langle \eta(\vec{r}, t) \rangle &= 0 \\ \langle \eta(\vec{r}, t) \eta(\vec{r}', t') \rangle &\sim \nabla^2 |\vec{r} - \vec{r}'|^{2\sigma-d} \delta(t - t'),\end{aligned}\tag{2}$$

where σ is an exponent characterizing the decay of spatial correlations (in the case of conservative white noise, which is of great interest in the MBE system, $\sigma = 0$).

After suggesting the model, Jung *et al.* [15, 16] investigated the NCKPZ problem using DRG analysis, and derived a very rich and complex picture of phases - each one of them described by a different set of critical exponents. In this letter I will not go into the details of all those phases, but would like to focus on the strong coupling behavior of NCKPZ. In this regime the DRG analysis yields the power counting result

$$\alpha = (2 - d - \rho + 2\sigma)/3, \quad z = (d + 10 - 2\rho - 2\sigma)/3,\tag{3}$$

where α is the roughness exponent and z is the dynamic exponent. This solution is said to be valid for positive values of ρ and for $d < 2 + 2\rho + 2\sigma$.

In this letter I will show that the above DRG analysis does not give the full description of the different phases in the strong coupling regime of NCKPZ. More specifically, I will show that an exact solution is possible for some one-dimensional cases (namely for $\rho = 2 + 2\sigma$). It is interesting to mention that this is the first time an exact result is suggested for the family of MBE processes. It will then be immediately seen that this exact solution is not compatible with the DRG result. Then, in order to clarify the situation, I will apply a method developed by Schwartz and Edwards [12, 22, 23] (also known as the Self-Consistent-Expansion (SCE) approach). This method has been previously applied successfully to the KPZ equation. The method gained much credit by being able to give a sensible prediction for the KPZ critical exponents in the strong coupling phase, where, as is well known, many Renormalization-Group (RG) approaches failed (as well as DRG of course) [24]. For the specific problem of NCKPZ, the SCE method indicates that two strong coupling solutions exist, one of which agrees with the power counting solution also obtained by DRG, while the other one recovers the exact one-dimensional result (and also extends it to higher dimensions). It turns out that each solution is valid for different values of the parameters σ , ρ and d (mutually excluding values).

In order to derive the exact result, I have to consider eqs. (1)-(2) in Fourier components

$$\frac{\partial h_q}{\partial t} = -Kq^4 h_q - \frac{1}{\sqrt{\Omega}} \sum_{\ell, m} \left(\lambda_0 + \lambda_\rho q^{-\rho} \right) q^2 \left(\vec{\ell} \cdot \vec{m} \right) \delta_{q, \ell+m} h_\ell h_m + \eta_q, \quad (4)$$

where η_q is the Fourier component of the noise that satisfies:

$$\begin{aligned} \langle \eta_q(t) \rangle &= 0 \\ \langle \eta_q(t) \eta_{q'}(t') \rangle &= 2D_0 q^{2-2\sigma} \delta_{q+q'} \delta(t-t'). \end{aligned} \quad (5)$$

I will show that for the case $\lambda_0 = 0$ an exact solution is possible for the steady state distribution resulting from eq. (4). However, before doing that I would like to mention that when $\rho > 0$, the λ_ρ term in eq. (4) dominates over the λ_0 term in the asymptotic small q limit, so that this term can be neglected. On the other hand when $\rho < 0$ the λ_ρ term can be neglected compared to the λ_0 term, and I retain the standard MBE results, unless $\lambda_0 = 0$ exactly (The specific case $\lambda_0 = 0$ and $\rho < 0$ is reasonable only under special physical conditions [25]). It will be understood therefore, that while the results obtained for positive ρ 's hold also for $\lambda_0 \neq 0$, the results obtained for negative ρ 's hold only when λ_0 is strictly zero. In other cases (i.e. $\rho < 0$ and $\lambda_0 \neq 0$) the MBE results are recovered.

I now turn from the NCKPZ equation in Langevin form to a Fokker-Planck form

$$\begin{aligned} \frac{\partial P}{\partial t} + \sum_q \frac{\partial}{\partial h_q} \left[D_0 q^{2-2\sigma} \frac{\partial}{\partial h_{-q}} + K q^4 h_q + \right. \\ \left. + \frac{\lambda_\rho q^{2-\rho}}{\sqrt{\Omega}} \sum_{\ell, m} \left(\vec{\ell} \cdot \vec{m} \right) \delta_{q, \ell+m} h_\ell h_m \right] P = 0, \end{aligned} \quad (6)$$

(where $P(\{h_q\}, t)$ is the probability of having the configuration $\{h_q\}$ at a specific time t).

Consider now the family of models defined by $\rho = 2 + 2\sigma$. By direct substitution, it can be verified immediately that the following probability function

$$P_0 = N \exp \left[-\frac{1}{2} \sum_q \frac{h_q h_{-q}}{\phi_q} \right], \quad (7)$$

(where N is the normalization yielding $\int P_0 \{h\} Dh = 1$, and ϕ_q is the two point correlation function) is a solution of the steady-state Fokker-Planck equation (i.e. with $\partial P / \partial t = 0$) for the one-dimensional case ($d = 1$) if $\phi_q = A q^{-\Gamma}$ with $A = D_0 / K$ and $\Gamma = \rho$. Translating this result to the frequently used notation in the field of surface growth I get

$$\alpha = \frac{\rho - 1}{2}, \quad (8)$$

where α is just the roughness exponent. This result is therefore consistent with the known exact solution of one-dimensional KPZ $\alpha = 1/2$ (by setting $\sigma = 0$ and $\rho = 2$). However, this result is manifestly different from the DRG result cited in eq. (3).

After establishing the exact result I turn to the application of the SCE method to NCKPZ in order to clarify the conflict as well as to complete the picture for any noise exponent σ and for higher dimensions. The SCE method is based on the Fokker-Planck form of NCKPZ equation, which is the reasonable thing to do once you have an exact solution for that equation in one dimension. The method constructs a self-consistent expansion of the distribution of the field concerned. The expansion is formulated in terms of ϕ_q and ω_q , where ϕ_q is the two-point function in momentum space, defined by $\phi_q = \langle h_q h_{-q} \rangle_S$, (the subscript S denotes steady state averaging), and ω_q is the characteristic frequency associated with h_q . I expect that for small enough q , ϕ_q and ω_q obey power laws in q ,

$$\phi_q = Aq^{-\Gamma} \quad \text{and} \quad \omega_q = Bq^\mu. \quad (9)$$

[Since dynamic surface growth is a remarkably multidisciplinary field, there are almost as many notations as there are workers in the field. Therefore I give a brief translation of our notations to those most frequently used:

$$z = \mu, \quad \alpha = (\Gamma - d)/2 \quad \text{and} \quad \beta = (\Gamma - d)/2\mu.] \quad (10)$$

The SCE method produces, to second order in this expansion, two nonlinear coupled integral equations in ϕ_q and ω_q , that can be solved exactly in the asymptotic small q limit to yield the required scaling exponents governing the steady state behavior and the time evolution.

By defining $M_{q\ell m} = \frac{\lambda_\rho}{\sqrt{\Omega}} q^{2-\rho} (\vec{\ell} \cdot \vec{m}) \delta_{q,\ell+m}$, $K_q = Kq^4$ and $D_q = D_0 q^{2-2\sigma}$ it can be seen that the NCKPZ equation is of the general form discussed in refs. [22, 23], where the SCE method is derived. Working to second order in the expansion, this yields the two coupled non-linear integral equations

$$D_q = K_q \phi_q + 2 \sum_{\ell, m} \frac{M_{q\ell m} M_{q\ell m} \phi_\ell \phi_m}{\omega_q + \omega_\ell + \omega_m} -$$

$$- 2 \sum_{\ell, m} \frac{M_{q\ell m} M_{\ell m q} \phi_m \phi_q}{\omega_q + \omega_\ell + \omega_m} - 2 \sum_{\ell, m} \frac{M_{q\ell m} M_{m\ell q} \phi_\ell \phi_q}{\omega_q + \omega_\ell + \omega_m} = 0, \quad (11)$$

and

$$K_q - \omega_q - 2 \sum_{\ell, m} M_{q\ell m} \frac{M_{\ell m q} \phi_m + M_{m\ell q} \phi_\ell}{\omega_\ell + \omega_m} = 0, \quad (12)$$

where in deriving the last equation I have used the Herring consistency equation [26]. In fact Herring's definition of ω_q is one of many possibilities, each leading to a different consistency equation. But it can be shown, as previously done in ref. [23], that this does not affect the exponents (universality).

A full solution of equations (11) and (12) in the limit of small q 's (i.e. large scales) yields a very rich family of solutions that will not interest us here, but may be the subject of a future detailed paper. Instead, I focus on the strong coupling solutions obtained by SCE. As mentioned above, two strong coupling solutions are obtained by SCE, each of them describes the critical exponents for a different set of the parameters ρ , σ and d .

The first solution is obtained just by power counting, and reads $\mu = (d + 10 - 2\rho - 2\sigma)/3$ and $\Gamma = (d + 4 - 2\rho + 4\sigma)/3$. This solution is valid for $\rho - 1 < \sigma$, $d < 6 - 2\sigma + \rho$, $d < 2 + 2\sigma + 2\rho$ and $d < 8 + 2\sigma - 4\rho$. Therefore, by direct inspection, this solution is not valid for the one-dimensional case with $\rho = 2 + 2\sigma$, and hence does not recover the exact Gaussian result obtained above. It should also be mentioned that when translated to frequently used notation (using eq. (10)) it can be seen that this solution is actually the same as the one obtained by the DRG analysis as in eq. (3).

The second solution is a non power counting solution. It turns out that for $d = 1$ eq. (11) is exactly solvable, and yields $\Gamma = \rho$. This result is actually the exact Gaussian solution of the one-dimensional case, mentioned above. In addition, from eq. (12), the dynamical exponent can be extracted for a certain range of the parameters, namely $1 < \rho < 3$ and $\rho > (4\sigma + 5)$, where it becomes $\mu = (9 - 3\rho)$.

For dimensionality higher than one (i.e. $d \geq 2$) such an exact solution in closed form cannot be found, and the strong coupling solution is determined from the combination of the scaling relation $\mu = (d + 8 - \Gamma - 2\rho)/2$, and the transcendental equation $F(\Gamma, \mu) = 0$, where F is given by

$$F(\Gamma, \mu) = - \int d^d t \frac{\vec{t} \cdot (\hat{e} - \vec{t})}{t^\mu + |\hat{e} - \vec{t}|^\mu + 1} \left[(\hat{e} \cdot \vec{t}) |\hat{e} - \vec{t}|^{2-\rho} t^{-\Gamma} + \hat{e} \cdot (\hat{e} - \vec{t}) t^{2-\rho} |\hat{e} - \vec{t}|^{-\Gamma} \right] +$$

$$+ \int d^d t \frac{[\vec{t} \cdot (\hat{e} - \vec{t})]^2}{t^\mu + |\hat{e} - \vec{t}|^\mu + 1} t^{-\Gamma} |\hat{e} - \vec{t}|^{-\Gamma}, \quad (13)$$

and $\hat{e}$ is a unit vector in an arbitrary direction. The last equation has to be solved numerically of course. In addition, This solution is valid as long as the solutions of the last equations satisfy the following four conditions: $d + 4 - 3\Gamma - 2\rho + 4\sigma < 0$, $d - 3\Gamma + 2\rho < 0$, $d < \Gamma$ and $d - 4 - \Gamma + 2\rho < 0$.

It should be mentioned that SCE yields many other "weak coupling" solutions to NCKPZ. They are called "weak coupling" in the sense that all of them yield the exponents of the linear MBE problem with correlated noise in the limit of $\rho = 0$. These solutions are also not obtained by DRG.

In summary, I have established an exact solution for the one-dimensional NCKPZ equation in the special case where $\rho = 2 + 2\sigma$. Then, I calculated the critical exponents for the most general case (i.e. any ρ , σ and d) using the SCE method, and obtained two possible strong coupling solutions. The first strong coupling solution is a power counting one, and is identical to the previous result of the DRG analysis. The second strong coupling solution is non power counting, and is not obtained by DRG. For $d = 1$, this solution turns out to be the exact Gaussian solution obtained at the beginning. This suggest that there is a big advantage in using the SCE to deal with such nonlinear problems.

Another advantage of using SCE has to do with the well known technical problem raised by Janssen [21]. Janssen suggested that there should be a correction to the scaling relation because of a nontrivial renormalization of the coupling constant. This correction appears in a two-loop RG calculation, but not in a one-loop calculation. Janssen himself suspected that in order to avoid this difficulty, a mode-coupling approach might be more appropriate. The SCE method is indeed closely related to the mode-coupling approaches, in the sense that similar (but not identical) equations are obtained, while the underlying derivation is different. Therefore, the fact SCE recovers the one-loop DRG power counting results of Jung *et al.*, for a specific range of the parameters [15, 16], is a nontrivial corroboration of that result.

Acknowledgement: I would like to thank Moshe Schwartz for useful discussions.

-
- [1] A.-L. Barabasi and H. E. Stanley, *Fractal Concepts in Surface Growth* (Cambridge Univ. Press, Cambridge, 1995).
- [2] M. Kardar, G. Parisi and Y.-C. Zhang, *Phys. Rev. Lett.* **56**, 889 (1986).
- [3] D. Wolf and J. Villain, *Europhys. Lett.* **13**, 389 (1990).
- [4] S. Das Sarma and P. Tamborenea, *Phys. Rev. Lett.* **66**, 325 (1991).
- [5] Z.-W. Lai and S. Das Sarma, *Phys. Rev. Lett.* **66**, 2348 (1991).
- [6] L.-H. Tang and T. Nattermann, *Phys. Rev. Lett.* **66**, 2899 (1991).
- [7] P.-M. Lam and F. Family, *Phys. Rev. A* **44**, 7939 (1991).
- [8] D. A. Kessler, H. Levine and L. M. Sander, *Phys. Rev. Lett.* **69**, 100 (1992).
- [9] T. Sun and M. Plischke, *Phys. Rev. Lett.* **71**, 3174 (1993).
- [10] E. Medina, T. Hwa, M. Kardar and Y.C. Zhang, *Phys. Rev. A* **39**, 3053 (1989).
- [11] A. Kr. Chattapohadhyay and J. K. Bhattacharjee, *Europhys. Lett.* **42**, 119 (1998).
- [12] E. Katzav and M. Schwartz, *Phys. Rev. E* **60**, 5677 (1999).
- [13] S. Mukherji and S. M. Bhattacharjee, *Phys. Rev. Lett.* **79**, 2502 (1997).
- [14] A. Kr. Chattapohadhyay, *Phys. Rev. E* **60**, 293 (1999).
- [15] Y. Jung, I.M. Kim and J.M. Kim, *Phys. Rev. E* **58**, 5467 (1998).
- [16] Y. Jung and I.M. Kim, *Phys. Rev. E* **62**, 2949 (2000).
- [17] G. Tang and B. Ma, *Int. Jour. Mod. Phys. B* **15**, 2275 (2001).
- [18] G. Tang and B. Ma, *Physica A* **298**, 257 (2001).
- [19] H. G. E. Hentschel and F. Family, *Phys. Rev. Lett.* **66**, 1982 (1991).
- [20] E. Katzav, *Physica A* **308**, 25 (2002).
- [21] H. K. Janssen, *Phys. Rev. Lett.* **78**, 1082 (1997).
- [22] M. Schwartz and S.F. Edwards, *Europhys. Lett.* **20**, 301 (1992).
- [23] M. Schwartz and S.F. Edwards, *Phys. Rev. E* **57**, 5730 (1998).
- [24] M. S. Li, T. Nattermann, H. Rieger and M. Schwartz, *Phys. Rev. B* **54**, 16024 (1996).
- [25] M. Schwartz, J. Villain, Y. Shapir and T. Nattermann, *Phys. Rev. B* **48**, 3095 (1993).
- [26] J.R. Herring, *Phys. Fluids* **8**, 2219 (1965); **9**, 2106 (1966).